Purbrook Junior School

MATHS
KNOWLEDGE ORGANISERS
YEAR 5

Number and Place Value   Knowledge Organiser

Key Vocabulary
millions
thousands
hundreds
tens
ones
zero
place value
greater than
less than
order
round
rounded
negative number
partition
digit
interval
sequence
linear sequence

Compare and Order

equals

26 + 38 = 8 × 8

Both calculations have the value 64.

greater than

23 873 > 8256

The number on the left has 2 ten thousands and the number on the right has 0 ten thousands.

less than

901 198 < 1 091 098

The number on the right has 1 million and the number on the left has 0 millions.

smallest | 898 | 6735 | 6835 | 7019 | 9002 | 11 235 | greatest

Negative Numbers

-25 -24 -23 -22 -21 -20 -19 -18 -17 -16 -15 -14 -13 -12 -11 -10 -9 -8 -7 -6 -5 -4 -3 -2 -1 0 1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24 25

Counting in Powers of 10

Counting in 10s

365	375	385	395	405	415

The tens increase until 9 tens becomes one more hundred and 0 tens.

Counting in 100s

2841	2941	3041	3141	3241	3341

The hundreds increase until 9 hundreds becomes one more thousand and 0 hundreds.

Counting in 10 000s

276 109	286 109	296 109	306 109

The ten thousands increase until 9 ten thousands become one more hundred thousand and 0 ten thousands.

Counting in 100 000s

2 972 151	3 072 151	3 172 151	3 272 151

The hundred thousands increase until 9 hundred thousands becomes one more million and 0 hundred thousands.

Number and Place Value

## Numbers to One Million

926 471

Hundred Thousands	Ten Thousands	Thousands	Hundreds	Tens	Ones
9	2	6	4	7	1

nine hundred and twenty-six thousand, four hundred and seventy-one

926 471 → 900 000, 20 000, 6000, 400, 70, 1

0   500 000   1 000 000

100 00   200 000   300 000   400 000   600 000   700 000   800 000   900 000

Roman Numerals

	I = 1	II = 2	III = 3	
IV = 4	V = 5	VI = 6	VII = 7	VIII = 8
IX = 9	X = 10	XI = 11	XX = 20	XXX = 30
XL = 40	L = 50	LX = 60	LXX = 70	LXXX = 80
XC = 90	C = 100	CL = 150	CC = 200	CCC = 300
CD = 400	D = 500	DC = 600	DCC = 700	DCCC = 800
CM = 900	M = 1000	MC = 1100	MD = 1500	MM = 2000

CCXLVIII = 248   DCCLXXXIV = 784   MMXIX = 2019

Rounding

Rounding to the nearest 10

20	21	22	23	24	25	26	27	28	29	30

← round down   round up →

Rounding to the nearest 1000

2000 ← 2499 | 2500 → 3000

round down   round up

Rounding to the nearest 100 000

200 000 ← 249 999 | 250 000 → 300 000

round down   round up

Addition and Subtraction — Knowledge Organiser

Key Vocabulary
Add
Total
Make
Plus
Sum
More
Altogether
Difference
Subtract
Less
Minus
Take away
Column addition
Column subtraction
Estimate
Inverse operation
Number facts
Place value
Complex

Addition

Place Value Grid: 3274 + 5601 = 8875

Column Method

Starting with the ones, add each column in turn. Regroup tens, hundreds, thousands, ten thousands and/or as required.

$$\begin{array}{r} 45864 \\ +23497 \\ \hline 69361 \\ \hline {\scriptstyle 1\,1\,1} \end{array}$$

Subtraction

Place Value Grid: 35 727 – 6313 = 29 414

TTh — 2 ten thousands left

Th — 5 thousands – 6 thousands cannot be done. Exchange ten thousand for ten thousands becoming 15 thousands – 6 thousands = 9 thousands

H — 7 hundreds – 3 hundreds = 4 hundreds

T — 2 tens – 1 ten = 1 ten

O — 7 ones – 3 ones = 4 ones

Column Method

Starting with the ones, subtract each column in turn. Exchange tens, hundreds, thousands and/or ten thousands as required.

$$\begin{array}{r} {\scriptstyle 6\ 13\ 1} \\ 35\not7\not42 \\ -\ 3476 \\ \hline 32266 \end{array}$$

Addition and Subtraction | Knowledge Organiser

Estimate and Approximate

Rounding to Estimate

$$41\ 635 + 7386 = 49\ 021$$

Round to ten:

$$41\ 630 + 7380 = 49\ 010$$

$$41\ 630 + 7390 = 49\ 020$$

$$41\ 640 + 7390 = 49\ 030$$

Rounding is not as accurate when both numbers are rounded up. A better estimate comes from "rounding" one down and one up.

Estimating on a Number Line

10 000   50 000

The arrow is about ¾ of the way across the line so it is 40 000.

Inverse Operations

Use the inverse to check:

53 476	
32 732	20 744

To check 53 476 – 32 732 = 20 744
use 32 732 + 20 744 = 53 476

Start with a number, subtract 409 and double. I end with 6264.
To find the starting number use the inverse: halve, then add 409. Half of 6264 = 3132. 3132 + 409 = 3541. The starting number was 3541.

Multistep Problems

Using a Bar Model
The sum of two numbers is 25 567.
The difference is 1875.

} 25 567
1875 }

Subtract 1875 from 25 567 = 23 692.
Halve 23 692 to find smaller number = 11 846.
Add 1875 to find larger number = 13 721.

£20		
£3.75	£8.49	?
£12.24		£7.76

£20 is used to buy 2 books costing £3.75 and £8.49.

How much change is given?

£3.75 + £8.49 = £12.24
£20.00 - £12.24 = £7.76

Multiplication and Division

Key Vocabulary

- multiply
- groups of
- lots of
- times
- divide
- share
- remainder
- factor
- multiple
- product

Factors

A factor is a number that divides into another number exactly, without leaving a remainder.

20

| 1 | 2 | 4 | | 5 | 10 | 20 |

A common factor is a factor of 2 or more numbers.

The factors of 20 are 1, 2, 4, 5, 10 and 20.

The factor pairs are:
1 and 20
2 and 10
4 and 5

Factors of 6
2
6
1 3
15
5
Factors of 15

Prime Numbers

1	2	3	4	5	6	7	8	9	10
11	12	13	14	15	16	17	18	19	20
21	22	23	24	25	26	27	28	29	30
31	32	33	34	35	36	37	38	39	40
41	42	43	44	45	46	47	48	49	50
51	52	53	54	55	56	57	58	59	60
61	62	63	64	65	66	67	68	69	70
71	72	73	74	75	76	77	78	79	80
81	82	83	84	85	86	87	88	89	90
91	92	93	94	95	96	97	98	99	100

Squared^2 and Cubed^3 Numbers

$2^2 = 4$
$2 \times 2 = 4$

$2^3 = 8$
$2 \times 2 \times 2 = 8$

$5^2 = 25$
$5 \times 5 = 25$

$5^3 = 125$
$5 \times 5 \times 5 = 125$

Related Calculations

$8 \times 9 = 72$
$80 \times 9 = 720$

$9 \times 8 = 72$
$90 \times 8 = 720$

$72 \div 9 = 8$
$720 \div 9 = 80$

$72 \div 8 = 9$
$720 \div 8 = 90$

Multiplication and Division

Short Multiplication

Revision

2543 × 7 = 17 801

	2	5	4	3
×				7
1	7	8	0	1
	1	3	3	2

Remember to move any regrouped digits into the next column. After the next multiplication, add the regrouped number to the answer.

Long Multiplication

2543 × 67 = 170 381

	2	5	4	3
×			6	7
1	7	8	0	1
	1	3	3	2
1	5	2	5	8	0
	1	3	2	1
1	7	0	3	8	1
		1		1

Before multiplying by the number in the tens column, remember to use zero as a placeholder because the 6 in 67 is 6 tens (60).

Division

136 ÷ 4 = 34

		3	4
4	1	3	6
−	1	2	0	→ 30 × 4
		1	6
	−	1	6	→ 4 × 4
			0

Short Division

		3	8
4	1	¹5	³2

15 ÷ 4 = 3 remainder 3

Remember to regroup any remainders and move them into the next column.

		4	5	5	r	3
5	2	2	²7	²8		

28 ÷ 5 = 5 remainder 3

If your calculation has a remainder, remember to record it in the answer using the letter **r**.

Decimals

Key Vocabulary

- tenths
- hundredths
- decimal tenths
- decimal hundredths
- decimal equivalents
- part-whole model
- rounding
- decimal point
- place value

Tenths, Hundredths and Thousandths

$\frac{0}{10}$ $\frac{1}{10}$ $\frac{2}{10}$ $\frac{3}{10}$ $\frac{4}{10}$ $\frac{5}{10}$ $\frac{6}{10}$ $\frac{7}{10}$ $\frac{8}{10}$ $\frac{9}{10}$ $\frac{10}{10}$

0 0.1 0.2 0.3 0.4 0.5 0.6 0.7 0.8 0.9 1

$\frac{0}{100}$ $\frac{1}{100}$ $\frac{2}{100}$ $\frac{3}{100}$ $\frac{4}{100}$ $\frac{5}{100}$ $\frac{6}{100}$ $\frac{7}{100}$ $\frac{8}{100}$ $\frac{9}{100}$ $\frac{1}{10}$

0 0.01 0.02 0.03 0.04 0.05 0.06 0.07 0.08 0.09 0.1

$\frac{0}{1000}$ $\frac{1}{1000}$ $\frac{2}{1000}$ $\frac{3}{1000}$ $\frac{4}{1000}$ $\frac{5}{1000}$ $\frac{6}{1000}$ $\frac{7}{1000}$ $\frac{8}{1000}$ $\frac{9}{1000}$ $\frac{1}{100}$

0 0.001 0.002 0.003 0.004 0.005 0.006 0.007 0.008 0.009 0.01

$\div 10$ $\div 10$ $\div 10$

$\frac{1}{1}$ $\frac{1}{10}$ $\frac{1}{100}$ $\frac{1}{1000}$

$\times 10$ $\times 10$ $\times 10$

Order and Compare Numbers with Three Decimal Places

Ones	Tenths	Hundredths	Thousandths
	$\frac{1}{10}$ $\frac{1}{10}$	$\frac{1}{100}$	$\frac{1}{1000}$ $\frac{1}{1000}$ $\frac{1}{1000}$

0 . 2 1 3

Ones	Tenths	Hundredths	Thousandths
1		$\frac{1}{100}$ $\frac{1}{100}$	$\frac{1}{1000}$ $\frac{1}{1000}$

1 . 0 2 2

Ones	Tenths	Hundredths	Thousandths
1 1	$\frac{1}{10}$		$\frac{1}{1000}$ $\frac{1}{1000}$ $\frac{1}{1000}$

2 . 1 0 3

Decimal Numbers as Fractions

$$0.71 = \frac{71}{100} = \frac{7}{10} + \frac{1}{100}$$

$$0.37 = \frac{37}{100} = \frac{3}{10} + \frac{7}{100}$$

Decimals

Multiplying and Dividing by 10, 100 and 1000

Tens	Ones	Tenths	Hundredths	Thousandths
3	8			
	3	8		
3	8			

÷ 10 / × 10

Tens	Ones	Tenths	Hundredths	Thousandths
3	8			
0	3	8		
3	8			

÷ 100 / × 100

Tens	Ones	Tenths	Hundredths	Thousandths
3	8			
0	0	3	8	
3	8			

÷ 1000 / × 1000

Adding and Subtracting Decimals

0.8 + 0.001 = 0.801
1.031 - 0.23 = 0.801
0.4005 + 0.4005 = 0.801

Rounding Decimals

1 ← 1.1 1.2 1.3 1.4 | 1.5 1.6 1.7 1.8 1.9 → 2

If the tenths digit is 1, 2, 3 or 4, we round down to the nearest whole number. | If the tenths digit is 5, 6, 7, 8 or 9, we round up to the nearest whole number.

1.1 ← 1.11 1.12 1.13 1.14 | 1.15 1.16 1.17 1.18 1.19 → 1.2

If the hundredths digit is 1, 2, 3 or 4, we round down to the nearest tenth. | If the hundredths digit is 5, 6, 7, 8 or 9, we round up to the nearest tenth.

Percentage and Decimal Equivalents

$50\% = \frac{50}{100} = \frac{1}{2} = 0.5$

$25\% = \frac{25}{100} = \frac{1}{4} = 0.25$

$10\% = \frac{10}{100} = \frac{1}{10} = 0.1$

$40\% = \frac{40}{100} = \frac{2}{5} = 0.4$

$20\% = \frac{20}{100} = \frac{1}{5} = 0.2$

$1\% = \frac{1}{100} = 0.01$

$70\% = \frac{70}{100} = \frac{7}{10} = 0.7$

Crossing the Whole

0.82 + 0.63 = 1.45
2.531 - 0.6 = 1.931

Fractions — Knowledge Organiser

Key Vocabulary

- numerator
- denominator
- unit fraction
- non-unit fraction
- whole
- equivalent
- mixed number
- improper fraction
- simplest form
- multiple
- common denominator
- common numerator

Equivalent Fractions

To find equivalent fractions, we multiply or divide the numerator and denominator by the same number.

$$\frac{1}{2} = \frac{5}{10} = \frac{50}{100}$$

(×5   ×10)

Compare and Order Fractions

We can compare and order fractions by using common denominators.

$$\frac{1}{3}, \frac{5}{6}, \frac{7}{12}$$

$$\frac{4}{12}, \frac{10}{12}, \frac{7}{12}$$

$$\frac{1}{3}, \frac{7}{12}, \frac{5}{6}$$

$$\frac{11}{8} > \frac{5}{4} = \frac{10}{8}$$

Mixed Numbers

Mixed numbers contain a whole number and a fraction.

whole → $2\frac{1}{4}$ ← fraction

Improper Fractions

An improper fraction has a numerator which is greater than or equal to the denominator.

$$\frac{5}{3}$$

Convert an Improper Fraction to a Mixed Number

$$\frac{9}{4}$$

$9 \div 4 = 2r1$ → $2\frac{1}{4}$

Divide the numerator by the denominator.

This shows you the whole number and the fraction.

Convert a Mixed Number to an Improper Fraction

Multiply the whole by the denominator to make an improper fraction.

$$2\frac{5}{6} = \frac{12}{6} + \frac{5}{6} = \frac{17}{6}$$

Add the fractions together.

Adding and Subtracting Fractions

To add or subtract fractions with denominators that are multiples of the same number, we must change one fraction to have the same denominator.

$$\frac{1}{3} + \frac{1}{3} = \frac{2}{3}$$

$$\frac{4}{5} - \frac{3}{5} = \frac{1}{5}$$

$$\frac{1}{4} + \frac{3}{8} = \frac{2}{8} + \frac{3}{8} = \frac{5}{8}$$

$$\frac{5}{6} - \frac{2}{3} = \frac{5}{6} - \frac{4}{6} = \frac{1}{6}$$

Fractions

Add Fractions Where the Total is Greater Than 1

$$\frac{1}{2} + \frac{3}{4} + \frac{5}{8} = \frac{4}{8} + \frac{6}{8} + \frac{5}{8} = \frac{15}{8} = 1\frac{7}{8}$$

Add Mixed Numbers

$$1\frac{1}{4} + \frac{3}{8} = 1\frac{2}{8} + \frac{3}{8} = 1 + \frac{5}{8} = 1\frac{5}{8}$$

$$1\frac{1}{4} + \frac{3}{8} = \frac{5}{4} + \frac{3}{8} = \frac{10}{8} + \frac{3}{8} = \frac{13}{8} = 1\frac{5}{8}$$

Multiply Unit Fractions by an Integer

$$\frac{1}{3} \times 5 = \frac{5}{3}$$

Multiply Non-Unit Fractions by an Integer

$$2 \times \frac{4}{9} = \frac{8}{9}$$

Multiply Mixed Numbers by Integers

Convert to an improper fraction and multiply the numerator by the integer.

$$2\frac{1}{4} \times 2 = \frac{9}{4} \times 2 = \frac{18}{4} = 4\frac{2}{4} = 4\frac{1}{2}$$

Use repeated addition.

$$2\frac{1}{4} \times 2 = 2\frac{1}{4} + 2\frac{1}{4} = 4\frac{2}{4} = 4\frac{1}{2}$$

Subtract from a Mixed Number

$$1\frac{2}{3} - \frac{2}{9} = 1\frac{6}{9} - \frac{2}{9} = 1\frac{4}{9}$$

starting number	find the equivalent fraction	subtract

Subtract Two Mixed Numbers

$$2\frac{3}{4} - 1\frac{5}{8} = 1\frac{1}{8}$$

$$2 - 1 = 1 \qquad \frac{3}{4} - \frac{5}{8} = \frac{1}{8}$$

Subtract from a Mixed Number - Breaking the Whole

$$2\frac{1}{4} - \frac{3}{8} = 2\frac{2}{8} - \frac{3}{8} = 1\frac{10}{8} - \frac{3}{8} = 1\frac{7}{8}$$

Converting Units

Key Vocabulary

mass
gram
kilogram
capacity
volume
millilitre
centilitre
litre
millimetre
centimetre
kilometre

Converting Mass

1000g = 1kg

$\frac{1}{10}$kg = 0.1kg = 100g

$\frac{1}{4}$kg = 0.25kg = 250g

$\frac{1}{2}$kg = 0.5kg = 500g

$\frac{3}{4}$kg = 0.75kg = 750g

kg ×1000 → g
kg ← ÷1000 g

Converting Capacity

1000ml = 1 litre

$\frac{1}{10}$l = 0.1l = 100ml

$\frac{1}{4}$l = 0.25l = 250ml

$\frac{1}{2}$l = 0.5l = 500ml

$\frac{3}{4}$l = 0.75l = 750ml

$\frac{1}{100}$l = 0.01l = 10ml

l ×1000 → ml
l ← ÷1000 ml

Converting Length

km ×1000 → m ×100 → cm ×10 → mm
km ← ÷1000 m ← ÷100 cm ← ÷10 mm

1000 metres = 1 kilometre

100cm = 1m

10mm = 1cm

$\frac{1}{10}$km = 0.1km = 100m

$\frac{1}{4}$km = 0.25km = 250m

$\frac{1}{2}$km = 0.5km = 500m

$\frac{3}{4}$km = 0.75km = 750m

Converting Units

Units of Time

Minute
1 minute = 60 seconds

Hour
1 hour = 60 minutes

Day
1 day = 24 hours

Week
1 week = 7 days

Monday
Tuesday
Wednesday
Thursday
Friday
Saturday
Sunday

Fortnight
1 fortnight = 2 weeks

Month
January = 31 days
February = 28 days (29 on a leap year)
March = 31 days
April = 30 days
May = 31 days
June = 30 days
July = 31 days
August = 31 days
September = 30 days
October = 31 days
November = 30 days
December = 31 days

Year
1 year =
12 months =
52 weeks =
365 days

Leap Year
1 leap year = 366 days

Decade
1 decade = 10 years

2000   2010

Century
1 century = 100 years

1900   2000

Millennium
1 millennium = 1000 years

1000   2000

Perimeter and Area

Knowledge Organiser

Key Vocabulary	Measure Perimeter	Calculate Perimeter

Key Vocabulary

- metre
- kilometre
- perimeter
- length
- width
- rectangle
- rectilinear
- dimensions

Measure Perimeter

Measure the perimeter of a rectangle:

w, l

Measure the length (l) and width (w).
Perimeter = l + w + l + w or (l + w) × 2

Measure the perimeter of regular shapes:

Measure the length (l) and count the number of sides (s) on the shape.

Perimeter = l × s

Measure the perimeter of irregular shapes:

Measure the length of each side and add them together.

Calculate Perimeter

Calculate the missing sides of this rectilinear shape to find the perimeter:

2cm
4cm
7cm
8cm
missing side 1
missing side 2

* This shape is not drawn to the dimensions specified.

Missing side 1 + 4cm = 8cm, so missing side 1 = 4cm.

Missing side 2 = 2cm + 7cm = 9cm

Perimeter = sum of all sides =
2cm + 4cm + 7cm + 4cm + 9cm + 8cm = 34cm

Length and Perimeter

Knowledge Organiser

Area of Rectangles

The area of a rectangle on a grid:

Multiply the length × width
= 6 × 3 = 18 squares.

The area of a rectangle = length (l) × width (w).

Area of Compound Shapes

To find the area of a compound shape, divide the shape into rectangles with known dimensions:

7cm

5cm

4cm   5cm

Area = 7cm × 4cm + 5cm × 5cm
     = 28cm² + 25cm²
     = 53cm²

Area of Irregular Shapes

To find the area of an irregular shape, find the number of whole squares and part squares.

Whole squares = 10
Part squares = 22

Estimate of area = whole squares + half part squares

= 10cm² + 11cm² = 21cm²

*There are other ways to estimate the area of irregular shapes.

Properties of Shape

Key Vocabulary

- angle
- right angle
- acute
- obtuse
- reflex
- protractor
- horizontal
- vertical
- parallel
- perpendicular
- polygon
- regular
- irregular
- two-dimensional
- three-dimensional
- flat face
- curved surface
- edge
- curved edge
- vertex
- apex

Regular and Irregular Polygons

Regular	Irregular

A polygon is any two-dimensional shape formed with straight lines.

In a regular polygon, all the sides and angles are equal.

In an irregular polygon, the sides and angles are not equal.

Representations

Cube models can be drawn as 2D representations using different elevations.

A shape net is a 2D drawing of an unfolded 3D shape. When you are drawing or reasoning about shape nets, think carefully about where the edges of the faces meet.

Shape net of a tetrahedron.

Properties of 3D Shapes

Name	Surfaces		Edges		Vertices	Picture
	Flat	Curved	Flat	Curved		
sphere	0	1	0	0	0	
cube	6	0	12	0	8	
cuboid	6	0	12	0	8	
cone	1	1	0	1	0	
cylinder	2	1	0	2	0	
square-based pyramid	5	0	8	0	5	
tetrahedron	4	0	6	0	4	
triangular prism	5	0	9	0	6	
pentagonal prism	7	0	15	0	10	
hexagonal prism	8	0	18	0	12	
octagonal prism	10	0	24	0	16	
octahedron	8	0	12	0	6	

A cone has an apex. This is because a vertex is the point where two straight edges meet and a cone has no straight edges.

Identifying Angles

Acute Angles
Any angle that measures less than 90° is called an **acute** angle.

Obtuse Angles
Any angle that measures greater than 90° and less than 180° is called an **obtuse** angle.

Reflex Angles
Any angle that measures greater than 180° is called a **reflex** angle.

117°
63°
Angles on a straight line always total 180°.

205°
65°
60°
30°
Angles around a point always total 360°.

Measuring and Drawing Angles
To measure angles, we use a protractor. Look carefully at how the numbers on the scale count from 0° to 180° in both directions.

Multiples of 90° can be used as descriptions of a turn.

$\frac{1}{4}$ turn = 90° $\frac{1}{2}$ turn = 180° $\frac{3}{4}$ turn = 270°   1 turn = 360°

Using Properties of Rectangles

6cm
7cm   2cm
6cm   8cm
13cm

6cm + 2cm = 8cm

7cm + 6cm = 13cm

Key Vocabulary

- coordinate
- quadrant
- x-axis
- y-axis
- reflection
- mirror line
- translation
- horizontal
- vertical

Coordinates are a useful way to locate a position on a map or grid.

The numbers across the horizontal line of the grid are on the **x-axis**.

The numbers on the vertical line of the grid are on the **y-axis**.

We always read or write the number on the x-axis before the y-axis.

The x and y position are written in brackets with a comma.

The coordinate of the orange spot is **(3, 4)**.

To help you remember which point to read or write first, simply remember to move 'along the corridor and up the stairs'.

In other words, move on the **x-axis** and then move on the **y-axis**.

Position and Direction

Reflection

A shape is reflected when is flipped over a mirror line.

The reflected image is congruent to the original. This means that the measurements of the sides and angles have not changed. Each point of the reflected shape is the same distance from the mirror line as the original shape.

Translation

In maths, translation means moving an object on a grid. The object is moved without changing the size, turning or reflecting it.

When translating an object on a grid, it can move up or down, left or right.

Volume

Key Vocabulary

- cubed
- area
- cross-section
- prism
- cube
- cuboid
- face
- length
- height
- width
- depth

Volume of Cubes and Cuboids

Volume is measured in cubed units. For example, **cm³**, **m³** and **km³**.

To calculate the volume of cubes and cuboids:

1. Calculate the area of the cross-section (one face).
2. Multiply the area of the cross-section (one face) by its depth.

Area of cross section (face) = 2cm × 2cm = 4cm²

4cm² × 2cm = Volume of 8cm³

Area of cross section (face) = 4cm × 2cm = 8cm²

8cm² × 3cm = Volume of 24cm³

Key Vocabulary

- axis
- continuous data
- horizontal
- data
- interpret
- label
- line graph
- maximum value
- minimum value
- pattern
- predict
- relationship
- represent
- scale
- survey
- table
- tally
- timetable
- vertical
- x-axis
- y-axis

Reading and Understanding Tables

A table to show ticket prices at a local cinema.

Ticket Type	Weekday Price	Weekend Price
Adult	£6	£7.50
Child	£4	£4.50
Student	£5.50	£6

In order to understand the data presented in a table, you must read the **table's title** and the **headings**. Remember to always look at the heading that **each piece of information** falls under.

Completing Tables

Here is a table showing the favourite drink flavours of some children.

	Boys	Girls	Total
Orange	8		18
Blackcurrant		6	
Total	15		

To find how many boys voted for blackcurrant, look at the total number of boys who voted and subtract the number of votes for orange.

To find how many girls voted for orange, look at the total number of votes for orange and subtract the number of votes from boys.

To find the total number of votes for blackcurrant, the total number of girls or the total number of voters, simply add up the values from the appropriate row or column.

Timetables

Here is a bus timetable:

Three different buses

	Mill Road	0726		0842
	High Street	0729	0803	
	Pitsmoor Road	0759	0833	
	Fulwood	0845	0919	0946

Bus stop locations

The bus starts at this time and location.

The bus does not stop here.

The bus terminates at this time and location.

Read and Interpret Line Graphs

Here is a line graph showing the average temperature for each month.

The y-axis shows temperature in intervals of 2°C on a scale of 0°C to 16°C.

The points show the average temperature for each month.

The x-axis shows the months of the year.

Use Line Graphs to Solve Problems

To find the average temperature in May, follow the arrow up from May and across to the temperature. As this is halfway between 10°C and 12°C, the average temperature in May is 11°C.

To find the difference between the average temperatures in August and in November, find the temperature for each month and calculate the difference between the two. The shape of the line graph can show how the temperature changed. he average temperature falls 9°C from August to November.

Draw Line Graphs

Here is a table showing the number of different types of fruit sold each day.

	Bananas	Apples
Mon	2	3
Tues	4	5
Wed	6	2
Thurs	5	4
Fri	8	1

This graph can be used to represent the data from the table.

Mark each point for the number of bananas sold each day and join each point with a line.

Mark each point for the number of apples sold each day and join each point with a line.